

Search for Anomalous $\gamma\gamma$ Production at TRISTAN

The AMY Collaboration

K.L. Sterner,^a A. Abashian,^a K. Gotow,^a D. Haim,^a M.E. Mattson,^a N. Morgan,^a L. Piilonen,^a P. Kirk,^b C.P. Cheng,^c W.X. Gao,^c W.G. Yan,^c M.H. Ye,^c S. Lusin,^d C. Rosenfeld,^d A.T.M. Wang,^d S. Wilson,^d L.Y. Zheng,^d C.A. Fry,^e R. Tanaka,^e R.E. Breedon,^{f,h} L.M. Chinitz,^f Winston Ko,^f R.L. Lander,^f J. Rowe,^f J.R. Smith,^f D. Stuart,^f S. Kanda,^g S.L. Olsen,^g K. Ueno,^g K. Abe,^h Y. Fujii,^h Y. Kurihara,^h F. Liu,^h A. Maki,^h T. Nozaki,^h T. Omori,^h H. Sagawa,^h Y. Sakai,^h T. Sasaki,^h Y. Sugimoto,^h Y. Takaiwa,^h S. Terada,^h R. Walker,^{h,n} F. Kajino,ⁱ R. Poling,^j T. Thomas,^j T. Aso,^k K. Miyano,^k H. Miyata,^k M. Oyoshi,^k M.H. Lee,^m F. Sannes,^m S. Schnetzer,^m R. Stone,^m J. Vinson,^m Y. Yamashita,^l A. Bodek,ⁿ B.J. Kim,ⁿ T. Kumita,ⁿ Y.K. Li,ⁿ C. Velisarris,ⁿ S. Kobayashi,^o A. Murakami,^o S.K. Sahu,^o Y.S. Chung,^p J.S. Kang,^p K.B. Lee,^p K.W. Park,^p S.K. Kim,^q S.S. Myung,^q S.K. Choi,^r D. Son,^r S. Ebara,^s S. Matsumoto,^s N. Takashimizu,^s and T. Ishizuka^t

^aVirginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA

^bLouisiana State University, Baton Rouge, LA 70803, USA

^cInstitute of High Energy Physics, Beijing 100039, China

^dUniversity of South Carolina, Columbia, SC 29208, USA

^eSSC Laboratory, Dallas, TX 75237, USA

^fUniversity of California, Davis, CA 95616, USA

^gUniversity of Hawaii, Honolulu, HI 96822, USA

^hKEK, National Laboratory for High Energy Physics, Ibaraki 305, Japan

ⁱKonan University, Kobe 658, Japan

^jUniversity of Minnesota, Minneapolis, MN 55455, USA

^kNiigata University, Niigata 950-21, Japan

^lNihon Dental College, Niigata 951, Japan

^mRutgers University, Piscataway, NJ 08854, USA

ⁿUniversity of Rochester, Rochester, NY 14627, USA

^oSaga University, Saga 840, Japan

^pKorea University, Seoul 136-701, South Korea

^qSeoul National University, Seoul 151-742, South Korea

^rKyungpook National University, Taegu 702-701, South Korea

^sChuo University, Tokyo 112, Japan

^tSaitama University, Urawa 338, Japan

Abstract

We report on measurements of the total cross section for $e^+e^- \rightarrow \gamma\gamma$ for center-of-mass energies between 57.4 and 59.5 GeV, using the AMY detector at the TRISTAN collider. We set new limits on the production of a possible new s -channel resonance decaying into photon pairs.

1. Introduction

Recently, the L3 Collaboration at LEP reported on an apparent excess of $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$ events ($\ell = \mu$ or e) in which the invariant mass of the photon pair was clustered around 59 GeV.^[1] The L3 Collaboration did not speculate on the origin of their events beyond noting that the probability of observing such clustering in any 5 GeV wide mass bin above 40 GeV was $\mathcal{O}(10^{-3})$, if the events were due purely to QED. The three other experiments at LEP have also reported on such events.^[2] Out of fifteen $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$ events found at LEP with $M_{\gamma\gamma} > 40$ GeV, five events were found with a $\gamma\gamma$ invariant mass in a 1 GeV range near 59 GeV.

A few speculative models have been advanced^[3,4] on the origin of the clustered events of L3 assuming that they are not due to QED. The models are based on the decay of the Z^0 into a massive object X and another boson, where X may be a scalar-pseudoscalar mixture or a spin-2 particle with a mass of approximately 60 GeV that decays predominantly into two photons. Regardless of the details of such models, if the particle X can couple with electrons, then the direct s -channel production may be observable in the TRISTAN energy range. If there is no coupling of X to fermions,

it can still be produced via an s -channel photon that results in $X \rightarrow \gamma\gamma$ plus a soft monochromatic photon. Previous studies of $e^+e^- \rightarrow \gamma\gamma, \gamma\gamma\gamma$ at TRISTAN^[5,6] were consistent with the QED prediction, however, no direct search for a narrow resonance was conducted.

We have searched for the direct production of a new state X via the reaction $e^+e^- \rightarrow X \rightarrow \gamma\gamma$, using the AMY detector at the TRISTAN e^+e^- collider, for center-of-mass energies $\sqrt{s}$ between 57.4 and 59.5 GeV in 250 MeV steps, with 1 to 2 pb⁻¹ of integrated luminosity per point (19 pb⁻¹ at 57.8 GeV).

2. The AMY Detector

The AMY detector is a general purpose solenoidal-type instrument employing two inner tracking chambers (VTX and ITC), a central drift chamber (CDC), and an electromagnetic calorimeter (SHC), all contained within a 3 Tesla magnetic field, and a barrel muon detector (MUO) outside the magnet return yoke. The end cap regions are instrumented with calorimeters (ESC) and small-angle luminosity monitors (SAC). The AMY detector has been described in detail elsewhere.^[7]

Final states with two or more photons are detected primarily by the SHC. This is a 14.4 radiation length gas-sampling calorimeter, consisting of twenty alternating layers of lead and resistive plastic proportional tubes.^[8] Each tube layer is sandwiched between cathode strips in the ϕ and θ directions. The SHC is constructed in six sextants, each covering an azimuthal range of 60° and a polar angle range of $|\cos\theta| < 0.73$.

Measurement of electromagnetic shower locations and energies in the SHC is de-

terminated by the signals from the cathode strips. The strips are arranged in projective towers that subtend an angle of 14.2 mr from the interaction point, and are ganged internally into five longitudinal divisions. The empirical energy resolution is found to be $\Delta E/E = 0.23/\sqrt{E} + 0.06$, with E in GeV. The large constant term is thought to be due to the effects of operating in the 3 T magnetic field. Shower centroids are typically located in ϕ and θ to less than one strip width.

Triggering

Within a given layer of the SHC, anode signals are ganged into 48 azimuthal sections; the ganged signals are summed externally over multiple layers to form trigger signals for 48 towers with either no segmentation or four longitudinal divisions. Three redundant combinations of these signals provide triggering for $e^+e^- \rightarrow \gamma\gamma$ events: (1) total energy trigger—the analog sum of all 48 trigger signals exceeding 8 GeV, (2) two-cluster trigger—two separate towers in coincidence with energies exceeding 5 GeV for one and 3 GeV for the other, and (3) shower shape trigger—coincidence between any two successive longitudinal divisions in a given tower, or in neighboring towers, with total energy exceeding 3 GeV.

Bhabha electrons are used to check the triggering efficiency independently, since they behave similarly to $\gamma\gamma$ events in the SHC. Bhabha events are triggered independently by the SHC triggers and by charged-track triggers derived from signals in the CDC and ITC. By examining the frequency of Bhabha events with a track trigger but no SHC trigger, we found the SHC trigger efficiency for Bhabha events to be essentially 100%. We assume, therefore, that the SHC triggering efficiency for $\gamma\gamma$ events exceeded 99% for all of the data of this study.

Luminosity Determination

To search for an anomaly in the production of $\gamma\gamma$ events, the most critical parameter aside from event statistics is the reliable determination of the integrated luminosity. This is accomplished in the AMY experiment by recording small-angle Bhabha events in the ESC, a sampling calorimeter constructed of alternating layers of lead and scintillator, with proportional tubes at the depth of electromagnetic shower maximum for position determination. The ESC covers the polar angle range $0.799 < |\cos\theta| < 0.982$. The systematic error in ESC luminosity is estimated to be $\sim 2\%$, and is dominated by the precision on alignment and fiducial definition.

3. Event Selection and Monte Carlo

The selection of $e^+e^- \rightarrow n\gamma$ events ($n \geq 2$) was performed according to the following criteria: (1) at least two SHC clusters with energies greater than $E_{beam}/3$ each, (2) the polar angle for each such cluster in the range $45^\circ < \theta < 135^\circ$, (3) the acollinearity angle between the two most energetic clusters of at most 10° , and (4) no charged tracks in the event. The resulting sample of 1054 events was visually scanned, resulting in the rejection of 19 background events from cosmic rays, misidentified Bhabhas, and SHC noise.

The cross section for $e^+e^- \rightarrow \gamma\gamma$, including radiative corrections to $\mathcal{O}(\alpha^3)$, was calculated by Monte Carlo integration using a program by Fujimoto, Igarashi, and Shimizu.^[9,10] This program is an implementation of the BASES/SPRING Monte Carlo package by Kawabata.^[11] The cross section of (34.81 ± 0.18) pb is based on a fully simulated sample of 1.06 fb^{-1} at 58 GeV that was subjected to the same selection

criteria as the experimental data. This value includes an event selection efficiency of $\varepsilon_{\gamma\gamma} = 0.933 \pm 0.011$.

We estimate the error in normalization for this analysis to be 2.3%, taking into account the uncertainty in the luminosity measurement and including the error in the $\gamma\gamma$ event selection efficiency.

4. Results

Table 1 summarizes the results of the energy scan. The center-of-mass energies $\sqrt{s}$ represent their actual values at the AMY interaction point rather than the nominal accelerator energies. The quantity σ_E represents the rms width of the center-of-mass energy distribution. The luminosities L are those obtained with the ESC. The quantity $N_{\gamma\gamma}$ [N_{QED}] represents the number of observed [simulated] $n\gamma$ ($n \geq 2$) events that pass the event selection criteria; N_{QED} is derived from the Monte Carlo simulation at 58 GeV discussed earlier, scaled by s and adjusted to the tabulated luminosity. The ratio of the observed $\gamma\gamma$ events to that expected from the Standard Model is also shown for each center-of-mass energy.

Barrel Bhabha events obtained concurrently with $\gamma\gamma$ events are used to test the internal consistency of the runs and to provide an additional check for systematic errors. For each center-of-mass energy, the ratio of observed barrel Bhabha events to ESC luminosity is scaled by s . These quantities are given in the last column of Table 1, and indicate the constancy in the operation of the SHC during the scan runs.

5. Analysis

The ratio of observed $n\gamma$ events ($n \geq 2$) to the QED expectation is plotted as a function of center-of-mass energy in Figure 1. The vertical error bars on the ratios are statistical only; the horizontal error bars indicate the rms spread in $\sqrt{s}$ rather than the error on the central values. A comparison of the data to QED alone, allowing the normalization to float without constraint, gives a fully consistent fit with $\chi^2/\text{dof} = 0.68$ for 8 degrees of freedom and a normalization value of 0.961 ± 0.029 . This fit is shown as the solid curve in Figure 1. The deviation of the normalization from unity is consistent, given the 2.9% statistical uncertainty from the fit on this parameter as well as the 2.3% normalization uncertainty mentioned above.

To examine the effect of a new state X , we again allow the normalization of the $\gamma\gamma$ data to float without constraint, and then compare these data to a model in which the $\gamma\gamma$ events are produced by an s -channel resonance X of mass M and total width Γ that sums incoherently with the conventional QED processes.

The effective cross section for production of X at center-of-mass energy $\sqrt{s}$, integrated over the solid angle of the SHC, is parameterized by a Breit-Wigner form:

$$\sigma_X(s) = (2J + 1)\Gamma_{\text{ee}}\text{BR}_{\gamma\gamma}f(s, M, \Gamma), \quad (1)$$

where

$$f(s, M, \Gamma) = (\Omega\varepsilon)_{\text{eff}} \frac{\pi}{s} \frac{\Gamma}{(M - \sqrt{s})^2 + \Gamma^2/4}. \quad (2)$$

Γ_{ee} is the partial width for $X \rightarrow e^+e^-$, $\text{BR}_{\gamma\gamma} \equiv \Gamma_{\gamma\gamma}/\Gamma$ is the branching ratio for $X \rightarrow \gamma\gamma$, and $(\Omega\varepsilon)_{\text{eff}}$ is the effective acceptance for detecting $e^+e^- \rightarrow \gamma\gamma$. We use

$(\Omega\varepsilon)_{\text{eff}} = \Omega_{\text{SHC}} \cdot \varepsilon_{\gamma\gamma} = 0.707 \cdot 0.933 = 0.659$. (This must be modified somewhat if the differential cross section is not isotropic.) The cross section σ_X is then convolved with the radiatively corrected beam resolution function^[12]

$$G_R(s, E) = \left(\frac{2\sigma_E}{\sqrt{s}} \right)^t \frac{t}{\sqrt{2\pi}\sigma_E} \int_0^\infty x^{t-1} \exp \left[-(z-x)^2/2 \right] dx, \quad (3)$$

where $z = (\sqrt{s} - E)/\sigma_E$, $t = 2(\alpha/\pi)[\ln(s/m_e^2) - 1]$, and σ_E is the width of the center-of-mass energy distribution from Table 1, to give the observed effective cross section

$$\tilde{\sigma}_X(s) = (2J+1)\Gamma_{\text{ee}}\text{BR}_{\gamma\gamma} \int_0^\infty f(E^2, M, \Gamma) G_R(s, E) dE. \quad (4)$$

The number of events expected from this process at center-of-mass energy $\sqrt{s}$ for an integrated luminosity L is given by $N_X(s) = \tilde{\sigma}_X(s) \cdot L$, and adds to the number expected from QED alone, $N_{QED}(s)$, listed in Table 1.

For a normalization A_o , the likelihood of observing $N_X(s_i)$ events from s -channel production of X at center-of-mass energy $\sqrt{s_i}$ is a rescaled Poisson distribution:^[13]

$$\begin{aligned} \mathcal{L}(s_i) = A_o \exp \{ N_{QED}(s_i) - A_o[N_X(s_i) + N_{QED}(s_i)] \} \\ \times \left\{ \frac{A_o[N_X(s_i) + N_{QED}(s_i)]}{N_{QED}(s_i)} \right\}^{N_{\gamma\gamma}(s_i)} \end{aligned} \quad (5)$$

and the overall likelihood is $\mathcal{L} = \prod_i \mathcal{L}(s_i)$.

This likelihood function peaks at $M = (58.35_{-0.08}^{+0.24})$ GeV, $\Gamma = (16 \pm 13)$ MeV and $(2J+1)\Gamma_{\text{ee}}\text{BR}_{\gamma\gamma} = (4.2_{-3.1}^{+3.4})$ keV, with a normalization of $A_o = 0.955 \pm 0.040$. The best-fit curve of $A_o(N_X + N_{QED})/N_{QED}$ is shown as the dashed curve in Figure 1. A comparison of this curve to the data has $\chi^2/dof = 0.47$ with 5 degrees of freedom.

Compared to $\chi^2/dof = 0.68$ with 8 degrees of freedom for QED alone, we conclude that this peak is not statistically significant.

We extract 90% confidence level upper limits on $(2J + 1)\Gamma_{ee}\text{BR}_{\gamma\gamma}$, for the normalization of 0.955 and particular choices of M and Γ , by numerical integration of the likelihood function.[†] These limits are shown in Figure 2. It is apparent that the limits are insensitive to the total width Γ when it is comparable to or smaller than the center-of-mass energy spread σ_E . Furthermore, the limits shift only slightly if the normalization is varied from its maximum likelihood value.

Upper limits on $(2J + 1)\Gamma_{ee}\text{BR}_{\gamma\gamma}$ at the 90% confidence level range from 0.5 keV to 8 keV, for an X mass between 57 and 59.6 GeV and a total width Γ below 100 MeV. Our limits can be compared directly with the expectation from the L3 observation of $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$ under the assumption that their clustered events arose from the production of a new state X . For example, if we assume that X is a scalar that couples only to photon and lepton pairs with a total width of $\Gamma \equiv \Gamma_{\gamma\gamma} + 3\Gamma_{ee}$, then an upper limit on $(2J + 1)\Gamma_{ee}\text{BR}_{\gamma\gamma}$ can be converted into an exclusion region in the plane of $\Gamma_{\gamma\gamma}$ vs. Γ_{ee} . Our 90% C.L. upper limit is shown in Figure 3 for a scalar X of mass 59 GeV.

Our limits may be incorporated with 90% C.L. limits of other experiments such as the L3 measurement^[1] of $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$ and the OPAL measurement^[16] of $e^+e^- \rightarrow \gamma\gamma\gamma$. For the former case, we calculate^[14] the cross section using the automatic amplitude generator GRACE^[15] including all possible diagrams. The limits from the three events with $M_{\gamma\gamma}$ clustered near 59 GeV observed by the L3 collaboration are

[†] Since the data are consistent with pure QED, the confidence *interval* includes the point $(2J + 1)\Gamma_{ee}\text{BR}_{\gamma\gamma} = 0$.

shown as the dashed curve in Figure 3, while the limits imposed by the absence of a $\ell^+\ell^-$ invariant mass near 59 GeV are given by the dash-dotted curve. The limits set by the non-observation of an excess of events with $M_{\gamma\gamma} = 59$ GeV in the OPAL $\gamma\gamma\gamma$ measurement are shown as the dotted curve of Figure 3. The limits of our measurement are thereby restricted to satisfying the condition that $\Gamma_{\gamma\gamma} < 10.6$ MeV and $\Gamma_{ee} < 120$ MeV. To summarize, the four *inclusion* regions overlap in the lower left of Figure 3.

For a (pseudo-)scalar that couples directly to pairs of fermions as well as photons, the exact forms of the decay widths for $X \rightarrow e^+e^-$ and $X \rightarrow \gamma\gamma$ are

$$\Gamma_{ee} = \frac{g_{ee}^2}{8\pi} M \quad (6)$$

and

$$\Gamma_{\gamma\gamma} = \frac{\alpha^2}{64\pi^3} \frac{M^3}{F_X^2} \quad (7)$$

respectively, where g_{ee} is the coupling constant of X to electrons, α is the fine structure constant, and F_X is a mass parameter analogous to f_π in π^0 decay. We use $\alpha/(\pi F_X)$ as the coupling constant of X to $\gamma\gamma$. Assuming that $\text{BR}_{\gamma\gamma} \sim 1$, we can convert the 90% C.L. upper limit on $\Gamma_{ee}\text{BR}_{\gamma\gamma}$ from our likelihood analysis into an upper limit on g_{ee} of 1.78×10^{-3} . We also place a 90% C.L. lower limit on F_X of 730 MeV that comes from the OPAL upper limit on $\Gamma_{\gamma\gamma}$ of 10.6 MeV.^[16] We examine the effect of these limits upon the $ee\gamma$ vertex correction value, $(g-2)_e$, which will have an additional contribution from this model of the form^[17]

$$\mathcal{A}_X \simeq \frac{m_e}{F_X} \frac{\alpha g_{ee}}{4\pi^3} \ln \frac{\Lambda}{M} \quad (8)$$

where Λ is the ultraviolet cutoff (fixed at 1 TeV). Our limits on F_X and g_{ee} im-

ply an upper bound of $\mathcal{A}_X < 2.0 \times 10^{-10}$, which is within the maximum allowed contribution^[18] to $(g-2)_e$ of 2.7×10^{-10} .

6. Conclusion

We have searched for resonant production of a new state X in e^+e^- collisions that couples to photon pairs at center-of-mass energies between 57.4 and 59.5 GeV. We find that the observed data are consistent with the QED prediction for $e^+e^- \rightarrow \gamma\gamma$ ($\chi^2/dof = 0.68$ for 8 degrees of freedom). Furthermore, we have extracted 90% C.L. upper limits on $(2J+1)\Gamma_{ee}\text{BR}_{\gamma\gamma}$ of 0.5—8.0 keV for the process $e^+e^- \rightarrow X \rightarrow \gamma\gamma$, should a state X exist with a mass between 57 and 59.6 GeV.

7. Acknowledgement

We thank the TRISTAN staff for the excellent operation of the storage ring. In addition we acknowledge the strong support provided by the staffs of our home institutions. We are particularly indebted to Professor Lay Nam Chang for fruitful discussions on the theoretical aspects of this paper. This work has been supported by the Japan Ministry of Education, Science and Culture (Monbusho), the Japan Society for the Promotion of Science, the U.S. Department of Energy, the U.S. National Science Foundation, the Korean Science and Engineering Foundation, the Ministry of Education of Korea and the Academia Sinica of the People's Republic of China.

REFERENCES

1. O. Adriani *et al.* (L3 Collaboration), Phys. Lett. **295B** (1992) 337.
2. S. Komamiya, K.W. Bell and S. Holmgren, private communications.
3. R. Garisto and J.N. Ng, Preprint TRI-PP-92-124, 16 Dec. 1992.
4. K. Hagiwara and S. Matsumoto, private communications.
5. K. Abe *et al.*, Z. Phys. C **45** (1989) 175.
6. K. Shimozawa *et al.*, Phys. Lett. **B284** (1992) 144.
7. AMY Collaboration, T. Kumita *et al.*, Phys. Rev. D **42** (1990) 1339.
8. A. Abashian *et al.*, to be published in Nucl. Instr. and Meth. A.
9. J. Fujimoto and M. Igarashi, Prog. Theor. Phys. **74** (1985) 791.
10. J. Fujimoto, M. Igarashi, Y. Shimizu, Prog. Theor. Phys. **77** (1987) 118.
11. S. Kawabata, Comp. Phys. Comm. **41** (1986) 127.
12. J.D. Jackson and D.L. Scharre, Nucl. Instr. and Meth. **128** (1975) 13.
13. O. Helene, Nucl. Instr. and Meth. **212** (1983) 319.
14. Y. Kurihara (unpublished).
15. H. Tanaka, Comp. Phys. Comm. **58** (1990) 153, H. Tanaka, T. Kaneko, and Y. Shimizu, Comp. Phys. Comm. **64** (1991) 149, T. Kaneko, in “New Computing Techniques in Physics Research,” ed. Perret-Gallix and W. Wojcik, (Edition du CRNS, Paris: 1990) 555.
16. M.Z. Akrawy *et al.* (OPAL Collaboration), Phys. Lett. **257B** (1991) 531.

17. F. del Águila, A. Méndez and R. Pascual, Phys. Lett. **140B** (1984) 431.
18. T. Kinoshita and W.B. Linquist, Phys. Rev. Lett. **47** (1981) 1573.

TABLE CAPTIONS

Table 1. Results of the energy scan. $\sqrt{s}$ is the center-of-mass energy, σ_E is the width of the center-of-mass energy distribution, L is the integrated luminosity from the ESC, N_{QED} is the number of events expected from QED alone, $N_{\gamma\gamma}$ is the number of $e^+e^- \rightarrow \gamma\gamma$ seen, and N_{BB}^*/L is the number of barrel Bhabha events seen per pb^{-1} of luminosity, scaled by s .

$\sqrt{s}$ (GeV)	σ_E (GeV)	L (pb ⁻¹)	N_{QED}	$N_{\gamma\gamma}$	$N_{\gamma\gamma}/N_{QED}$	N_{BB}^*/L
57.374	0.095	2.05	72.9 ± 0.4	69	0.95 ± 0.11	265 ± 11
57.772	0.096	19.55	680.5 ± 3.4	636	0.935 ± 0.037	276.0 ± 3.8
57.972	0.096	1.39	48.4 ± 0.2	49	1.01 ± 0.15	272 ± 14
58.220	0.098	1.33	45.8 ± 0.2	50	1.09 ± 0.15	259 ± 14
58.470	0.098	1.33	45.6 ± 0.2	54	1.18 ± 0.16	308 ± 15
58.718	0.099	1.73	58.7 ± 0.3	61	1.04 ± 0.13	278 ± 13
58.968	0.100	1.36	45.9 ± 0.2	45	0.98 ± 0.15	284 ± 15
59.216	0.100	1.21	40.5 ± 0.2	45	1.11 ± 0.17	294 ± 16
59.466	0.102	0.97	32.3 ± 0.2	26	0.81 ± 0.16	236 ± 16

FIGURE CAPTIONS

Figure 1. Ratio of observed $\gamma\gamma$ events to the QED prediction. The solid curve is the QED prediction, with $\chi^2/dof = 0.68$ for 8 degrees of freedom and a normalization of 0.961. The dashed curve is the maximum likelihood prediction, with $\chi^2/dof = 0.47$ for 5 degrees of freedom and a normalization of 0.955.

Figure 2. 90% C.L. upper limits on $(2J+1)\Gamma_{ee}\text{BR}_{\gamma\gamma}$ as a function of mass M of the state X , for widths of 1 MeV (solid curve), 100 MeV (dashed curve), 1 GeV (dotted curve) and 3 GeV (dash-dotted curve), with a normalization of 0.955. The region above a given curve is excluded.

Figure 3. Exclusion regions (90% C.L.) on $\Gamma_{\gamma\gamma}$ *vs.* Γ_{ee} for a scalar X with $M \sim 59$ GeV, assuming $\Gamma \equiv \Gamma_{\gamma\gamma} + 3\Gamma_{ee}$. The region above and to the right of a given curve is excluded. The solid curve is from this experiment. The other curves are from the observation of up to three $\ell^+\ell^-\gamma\gamma$ events with $M_{\gamma\gamma} \sim 59$ GeV (dashed) and the absence of $\ell^+\ell^-\gamma\gamma$ events with $M_{\ell^+\ell^-} \sim 59$ GeV (dash-dotted) (Ref. 1), and from the lack of a signal at $M_{\gamma\gamma} \sim 59$ GeV in $\gamma\gamma\gamma$ events (dotted) (Ref. 16).